Neuroscience Letters, 148 (1992) 216–218
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Abe, H. and Rusak, B., Stimulation of the hamster ventral lateral geniculate nucleus induces Fos-like immunoreactivity in suprachiasmatic nucleus cells, 185
Abe, T., see Shimosegawa, T., 67
Ábrahám, C.S., see Megyeri, P., 137
Aijón, J., see Alonso, J.R., 101
Akaike, N., see Takahashi, K., 60
Akiyama, H., see Ikeda, K., 126
Albala, N., see Mailleux, P., 177
Alonso, J.R., Sánchez, F., Arévalo, R., Carretero, J., Vázquez, R. and Aijón, J., Partial coexistence of NADPH-diaphorase and somatostatin in the rat hypothalamic paraventricular nucleus, 101
Arendt, T. and Brückner, M.K., Perisomatic sprouts immunoreactive for nerve growth factor receptor and neurofibrillary degeneration affect different neuronal populations in the basal nucleus in patients with Alzheimer's disease, 63
Arévalo, R., see Alonso, J.R., 101
Arioka, M., see Ishiguro, K., 202
Asakura, T., see Shimosegawa, T., 67
Avoli, M., see Siniscalchi, A., 159

Ballanyi, K., Kuwana, S.-i., Völker, A., Morawietz, G. and Richter, D.W., Developmental changes in the hypoxia tolerance of the in vitro respiratory network of rats, 141
Bell, C., see Hendry, I.A., 117
Bertrand, N., Ishii, H. and Spats, M., Regional and temporal glycerol changes induced by forebrain ischemia in gerbils, 81
Bianchi, M., Sacerdote, P., Ricciardi-Castagnoli, P., Mantegazza, P. and Panerai, A.E., Central effects of tumor necrosis factor α and interleukin-1α on nociceptive thresholds and spontaneous locomotor activity, 76
Bitran, M., see Torres, G., 39
Bloedel, J.R., see Irwin, K.B., 71
Bornstein, J.C., see Costa, M., 121
Bracha, V., see Irwin, K.B., 71
Bredt, D.S., see Costa, M., 121
Brilliant, M., Elble, R.J., Ghobrial, M. and Struble, R.G., Distribution of amyloid in the brainstem of patients with Alzheimer disease, 23
Brookes, S.J.H., see Costa, M., 121
Brückner, M.K., see Arendt, T., 63

Cano, J., see Santiago, M., 145
Carretero, J., see Alonso, J.R., 101
Carter Snead III, C., Evidence for G protein modulation of experimental-generalized absence seizures in rat, 15
Chan, J.Y.H., see Shih, C.-D., 169
Chan, S.H.H., see Shih, C.-D., 169
Costa, M., Furness, J.B., Pompolo, S., Brookes, S.J.H., Bornstein, J.C., Bredt, D.S. and Snyder, S.H., Projections and chemical coding of neurons with immunoreactivity for nitric oxide synthase in the guinea-pig small intestine, 121
Craig, A.D., see Irwin, K.B., 71
Cuadros, M.A., Moujahid, A., Martin-Partido, G. and Navascués, J., Microglia in the mature and developing quail brain as revealed by a monoclonal antibody recognizing hemopoietic cells, 11

da Silva, F.H.L., see Kamphuis, W., 51
Daelmans, H.E.M., see Witter, M.P., 164
De Rijk, T.C., see Kamphuis, W., 51
Do, M.C. and Gilles, M., Effects of reducing plantar support on anticipatory postural and intentional activities associated with flexion of the lower limb, 181

Ebihara, S., see Hasegawa, M., 89
Elble, R.J., see Brilliant, M., 23
Erneux, C., see Mailleux, P., 177

Feuerstein, G., see Lysko, P.G., 34
Florack, V.J., see Watt, C.B., 47
Fujii, S., Wakizaka, A., Sekino, Y., Kuroda, Y., Ito, K.-i., Miyakawa, H. and Kato, H., Adenosine A_2 receptor antagonist facilitates the reversal of long-term potentiation (depotentiation) of evoked postsynaptic potentials but inhibits that of population spikes in hippocampal CA1 neurons, 148
Furness, J.B., see Costa, M., 121

Geertsen, S., Trifaro, J.-M. and Quik, M., Phorbol esters and K^+ upregulate α-bungarotoxin binding sites in cultured chromaffin cells through a related mechanism, 207
Ghobrial, M., see Brilliant, M., 23
Gilles, M., see Do, M.C., 181
Greene, R.W., see Luebke, J.I., 129
Guilarte, T.R. and Miceli, R.C., Age-dependent effects of lead on [^{3}H]MK-801 binding to the NMDA receptor-gated ionophore: in vitro and in vivo studies, 27

Haga, C., see Ikeda, K., 126
Haley, J.E., see Kitto, K.F., 1
Han, Z.-S. and Ju, G., Responses of neurons in the rat anterior hypothalamic periventricular zone to osmotic stimulation and angiotensin II in vitro, 190
Hasegawa, M. and Ebihara, S., Circadian rhythms of pineal melatonin release in the pigeon measured by in vivo microdialysis, 89
Hendry, I.A., Messina, A. and Bell, C., Neonatal nerve growth factor treatment alters the preganglionic innervation pattern of rat superior cervical ganglion, 117
Herrera-Marschitz, M., see Morino, P., 133
Hökfelt, T., see Morino, P., 133
Hösli, E., see Hösli, L., 114
Hösli, L., Hösli, E., Kaeser, H. and Lefkovits, M., Colocalization of receptors for vasoactive peptides on astrocytes of cultured rat spinal cord and brain stem: electrophysiological effects of atrial and brain natriuretic peptide, neuropeptide Y and bradykinin, 114
Huidobro-Toro, J.P., see Torres, G., 39

Iizuka, M., see Kiehn, O., 43
Ikeda, K., Haga, C., Akiyama, H., Kase, K. and Iritani, S., Coexistence of paired helical filaments and glial filaments in astrocytic processes within ghost tangles, 126
Imahori, K., see Ishiguro, K., 202
Iritani, S., see Ikeda, K., 126
Irwin, K.B., Craig, A.D., Bracha, V. and Bloedel, J.R., Distribution of c-*fos* expression in brainstem neurons associated with conditioning and pseudo-condititioning of the rabbit nictitating membrane reflex, 71
Ishiguro, K., Omori, A., Takamatsu, M., Sato, K., Arioka, M.,

Neuroscience Letters, 148 (1992) 216–218
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Abe, H. and Rusak, B., Stimulation of the hamster ventral lateral geniculate nucleus induces Fos-like immunoreactivity in suprachiasmatic nucleus cells, 185
Abe, T., see Shimosegawa, T., 67
Ábrahám, C.S., see Megyeri, P., 137
Aijón, J., see Alonso, J.R., 101
Akaike, N., see Takahashi, K., 60
Akiyama, H., see Ikeda, K., 126
Albala, N., see Mailleux, P., 177
Alonso, J.R., Sánchez, F., Arévalo, R., Carretero, J., Vázquez, R. and Aijón, J., Partial coexistence of NADPH-diaphorase and somatostatin in the rat hypothalamic paraventricular nucleus, 101
Arendt, T. and Brückner, M.K., Perisomatic sprouts immunoreactive for nerve growth factor receptor and neurofibrillary degeneration affect different neuronal populations in the basal nucleus in patients with Alzheimer's disease, 63
Arévalo, R., see Alonso, J.R., 101
Arioka, M., see Ishiguro, K., 202
Asakura, T., see Shimosegawa, T., 67
Avoli, M., see Siniscalchi, A., 159

Ballanyi, K., Kuwana, S.-i., Völker, A., Morawietz, G. and Richter, D.W., Developmental changes in the hypoxia tolerance of the in vitro respiratory network of rats, 141
Bell, C., see Hendry, I.A., 117
Bertrand, N., Ishii, H. and Spats, M., Regional and temporal glycerol changes induced by forebrain ischemia in gerbils, 81
Bianchi, M., Sacerdote, P., Ricciardi-Castagnoli, P., Mantegazza, P. and Panerai, A.E., Central effects of tumor necrosis factor α and interleukin-1α on nociceptive thresholds and spontaneous locomotor activity, 76
Bitran, M., see Torres, G., 39
Bloedel, J.R., see Irwin, K.B., 71
Bornstein, J.C., see Costa, M., 121
Bracha, V., see Irwin, K.B., 71
Bredt, D.S., see Costa, M., 121
Brilliant, M., Elble, R.J., Ghobrial, M. and Struble, R.G., Distribution of amyloid in the brainstem of patients with Alzheimer disease, 23
Brookes, S.J.H., see Costa, M., 121
Brückner, M.K., see Arendt, T., 63

Cano, J., see Santiago, M., 145
Carretero, J., see Alonso, J.R., 101
Carter Snead III, C., Evidence for G protein modulation of experimental-generalized absence seizures in rat, 15
Chan, J.Y.H., see Shih, C.-D., 169
Chan, S.H.H., see Shih, C.-D., 169
Costa, M., Furness, J.B., Pompolo, S., Brookes, S.J.H., Bornstein, J.C., Bredt, D.S. and Snyder, S.H., Projections and chemical coding of neurons with immunoreactivity for nitric oxide synthase in the guinea-pig small intestine, 121
Craig, A.D., see Irwin, K.B., 71
Cuadros, M.A., Moujahid, A., Martin-Partido, G. and Navascués, J., Microglia in the mature and developing quail brain as revealed by a monoclonal antibody recognizing hemopoietic cells, 11

da Silva, F.H.L., see Kamphuis, W., 51
Daelmans, H.E.M., see Witter, M.P., 164
De Rijk, T.C., see Kamphuis, W., 51
Do, M.C. and Gilles, M., Effects of reducing plantar support on anticipatory postural and intentional activities associated with flexion of the lower limb, 181

Ebihara, S., see Hasegawa, M., 89
Elble, R.J., see Brilliant, M., 23
Erneux, C., see Mailleux, P., 177

Feuerstein, G., see Lysko, P.G., 34
Florack, V.J., see Watt, C.B., 47
Fujii, S., Wakizaka, A., Sekino, Y., Kuroda, Y., Ito, K.-i., Miyakawa, H. and Kato, H., Adenosine A_2 receptor antagonist facilitates the reversal of long-term potentiation (depotentiation) of evoked postsynaptic potentials but inhibits that of population spikes in hippocampal CA1 neurons, 148
Furness, J.B., see Costa, M., 121

Geertsen, S., Trifaro, J.-M. and Quik, M., Phorbol esters and K^+ upregulate α-bungarotoxin binding sites in cultured chromaffin cells through a related mechanism, 207
Ghobrial, M., see Brilliant, M., 23
Gilles, M., see Do, M.C., 181
Greene, R.W., see Luebke, J.I., 129
Guilarte, T.R. and Miceli, R.C., Age-dependent effects of lead on [^{3}H]MK-801 binding to the NMDA receptor-gated ionophore: in vitro and in vivo studies, 27

Haga, C., see Ikeda, K., 126
Haley, J.E., see Kitto, K.F., 1
Han, Z.-S. and Ju, G., Responses of neurons in the rat anterior hypothalamic periventricular zone to osmotic stimulation and angiotensin II in vitro, 190
Hasegawa, M. and Ebihara, S., Circadian rhythms of pineal melatonin release in the pigeon measured by in vivo microdialysis, 89
Hendry, I.A., Messina, A. and Bell, C., Neonatal nerve growth factor treatment alters the preganglionic innervation pattern of rat superior cervical ganglion, 117
Herrera-Marschitz, M., see Morino, P., 133
Hökfelt, T., see Morino, P., 133
Hösli, E., see Hösli, L., 114
Hösli, L., Hösli, E., Kaeser, H. and Lefkovits, M., Colocalization of receptors for vasoactive peptides on astrocytes of cultured rat spinal cord and brain stem: electrophysiological effects of atrial and brain natriuretic peptide, neuropeptide Y and bradykinin, 114
Huidobro-Toro, J.P., see Torres, G., 39

Iizuka, M., see Kiehn, O., 43
Ikeda, K., Haga, C., Akiyama, H., Kase, K. and Iritani, S., Coexistence of paired helical filaments and glial filaments in astrocytic processes within ghost tangles, 126
Imahori, K., see Ishiguro, K., 202
Iritani, S., see Ikeda, K., 126
Irwin, K.B., Craig, A.D., Bracha, V. and Bloedel, J.R., Distribution of c-*fos* expression in brainstem neurons associated with conditioning and pseudo-condititioning of the rabbit nictitating membrane reflex, 71
Ishiguro, K., Omori, A., Takamatsu, M., Sato, K., Arioka, M.,

Uchida, T. and Imahori, K., Phosphorylation sites on tau by tau protein kinase I, a bovine derived kinase generating an epitope of paired helical filaments, 202
Ishii, H., see Bertrand, N., 81
Ito, K.-i., see Fujii, S., 148
Ito, S.-i., Multiple projection of vagal nonmyelinated afferents to the anterior insular cortex in rats, 151

Jorritsma-Byham, B., see Witter, M.P., 164
Ju, G., see Han, Z.-S., 190

Kaeser, H., see Hösli, L., 114
Kameda, H., see Takahashi, K., 60
Kamphuis, W., Monyer, H., De Rijk, T.C. and da Silva, F.H.L., Hippocampal kindling increases the expression of glutamate receptor-A Flip and -B Flip mRNA in dentate granule cells, 51
Kase, K., see Ikeda, K., 126
Kataoka, M., see Takahashi, K., 60
Kato, H., see Fujii, S., 148
Kawano, H. and Masuko, S., Met-enkephalin-Arg^6-Gly^7-Leu^8- and substance P-containing projections from the nucleus preopticus medianus to the paraventricular hypothalamic nucleus, 211
Kiehn, O., Iizuka, M. and Kudo, N., Resetting from low threshold afferents of *N*-methyl-D-aspartate-induced locomotor rhythm in the isolated spinal cord-hindlimb preparation from newborn rats, 43
Kitto, K.F., Haley, J.E. and Wilcox, G.L., Involvement of nitric oxide in spinally mediated hyperalgesia in the mouse, 1
Koizumi, M., see Shimosegawa, T., 67
Kondo, Y., see Tomura, H., 93
Kovács, J., see Megyeri, P., 137
Koyama, I., see Sekino, Y., 109
Krause, J.E., see Marksteiner, J., 55
Kudo, N., see Kiehn, O., 43
Kuroda, Y., see Fujii, S., 148
Kuwana, S.-i., see Ballanyi, K., 141

Larsen, P.J., see Mikkelsen, J.D., 195
Lefkovits, M., see Hösli, L., 114
Luebke, J.I., Weider, J.M., McCarley, R.W. and Greene, R.W., Distribution of NADPH-diaphorase positive somata in the brainstem of the monitor lizard *Varanus exanthematicus*, 129
Lysko, K.A., see Lysko, P.G., 34
Lysko, P.G., Lysko, K.A., Webb, C.L. and Feuerstein, G., Neuroprotective effects of carvedilol, a new antihypertensive, at the *N*-methyl-D-aspartate receptor, 34

Machado, A., see Santiago, M., 145
Mailleux, P., Takazawa, K., Albala, N., Erneux, C. and Vanderhaeghen, J.-J., Astrocytic localization of the messenger RNA encoding the isoenzyme B of inositol (1,4,5) 3-kinase in the human brain, 177
Mailleux, P. and Vanderhaeghen, J.-J., Localisation of cannabinoid receptor in the human developing and adult basal ganglia. Higher levels in the striatonigral neurons, 173
Mantegazza, P., see Bianchi, M., 76
Marksteiner, J., Saria, A., Miller, C.H. and Krause, J.E., Differential increases of neurokinin B- and enkephalin-like immunoreactivities and their mRNAs after chronic haloperidol treatment in the rat, 55
Martin-Partido, G., see Cuadros, M.A., 11
Masuko, S., see Kawano, H., 211
McAdoo, D.J., see Paleckova, V., 19
McCarley, R.W., see Luebke, J.I., 129
McGeer, E.G., Singh, E.A. and McGeer, P.L., Apparent anterograde transport of basic fibroblast growth factor in the rat nigrostriatal dopamine system, 31
McGeer, P.L., see McGeer, E.G., 31
Meana, J.J., see Morino, P., 133
Megyeri, P., Ábrahám, C.S., Temesvári, P., Kovács, J., Vas, T. and Speer, C.P., Recombinant human tumor necrosis factor α constricts pial arterioles and increases blood–brain permeability in newborn piglets, 137
Messina, A., see Hendry, I.A., 117
Miceli, R.C., see Guilarte, T.R., 27
Mikkelsen, J.D. and Larsen, P.J., A high concentration of NPY (Y_1)-receptor mRNA-expressing cells in the rat arcuate nucleus, 195
Miller, C.H., see Marksteiner, J., 55
Miyakawa, H., see Fujii, S., 148
Moerlein, S.M. and Perlmutter, J.S., Specific binding of 3*N*-(2'-[^{18}F]fluoroethyl)benperidol to primate cerebral dopaminergic D_2 receptors demonstrated in vivo by PET, 97
Monyer, H., see Kamphuis, W., 51
Morawietz, G., see Ballanyi, K., 141
Morino, P., Herrera-Marschitz, M., Meana, J.J., Ungerstedt, U. and Hökfelt, T., Immunohistochemical evidence for a crossed cholecystokinin corticostriatal pathway in the rat, 133
Moujahid, A., see Cuadros, M.A., 11
Murthy, C., see Ratnakumari, L., 85

Nakai, Y., see Shioda, S., 155
Navascués, J., see Cuadros, M.A., 11

O'Callaghan, J.P. and Seidler, F.J., 1-Methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP)-induced astrogliosis does not require activation of ornithine decarboxylase, 105
Okajima, F., see Tomura, H., 93
Omori, A., see Ishiguro, K., 202

Palecek, J., see Paleckova, V., 19
Paleckova, V., Palecek, J., McAdoo, D.J. and Willis, W.D., The non-NMDA antagonist CNQX prevents release of amino acids into the rat spinal cord dorsal horn evoked by sciatic nerve stimulation, 19
Panerai, A.E., see Bianchi, M., 76
Perlmutter, J.S., see Moerlein, S.M., 97
Pompolo, S., see Costa, M., 121
Powel, R.J., see Pullan, L.M., 199
Pullan, L.M. and Powel, R.J., Comparison of binding at strychnine-sensitive (inhibitory glycine receptor) and strychnine-insensitive (*N*-methyl-D-aspartate receptor) glycine binding sites, 199

Quik, M., see Geertsen, S., 207

Ratnakumari, L. and Murthy, C., In vitro and in vivo effects of ammonia on glucose metabolism in the astrocytes of rat cerebral cortex, 85
Ricciardi-Castagnoli, P., see Bianchi, M., 76
Richter, D.W., see Ballanyi, K., 141
Rusak, B., see Abe, H., 185
Rustioni, A., see Valtschanoff, J.G., 6

Sacerdote, P., see Bianchi, M., 76
Sánchez, F., see Alonso, J.R., 101
Santiago, M., Machado, A. and Cano, J., Fast sodium channel dependency of the somatodendritic release of dopamine in the rat's brain, 145
Saria, A., see Marksteiner, J., 55

Sato, K., see Ishiguro, K., 202
Satoh, A., see Shimosegawa, T., 67
Schmidt, H.H.H.W., see Valtschanoff, J.G., 6
Seidler, F.J., see O'Callaghan, J.P., 105
Sekino, Y., see Fujii, S., 148
Sekino, Y. and Koyama, I., Selective inhibition of homosynaptic depression in a tetanized pathway by an adenosine A_1 blocker in the CA1 region of rat hippocampal slice, 109
Shih, C.-D., Chan, J.Y.H. and Chan, S.H.H., Tonic suppression of baroreceptor reflex response by endogenous neuropeptide Y at the nucleus tractus solitarius of the rat, 169
Shimoda, Y., see Shioda, S., 155
Shimosegawa, T., Abe, T., Satoh, A., Asakura, T., Yoshida, K., Koizumi, M. and Toyota, T., Histochemical demonstration of NADPH-diaphorase activity, a marker for nitric oxide synthase, in neurons of the rat pancreas, 67
Shioda, S., Shimoda, Y. and Nakai, Y., Ultrastructural studies of medullary synaptic inputs to vasopressin-immunoreactive neurons in the supraoptic nucleus of the rat hypothalamus, 155
Singh, E.A., see McGeer, E.G., 31
Siniscalchi, A. and Avoli, M., Modulation by $GABA_B$ receptors of spontaneous synchronous activities induced by 4-aminopyridine in the rat hippocampus, 159
Snyder, S.H., see Costa, M., 121
Spats, M., see Bertrand, N., 81
Speer, C.P., see Megyeri, P., 137
Staiger, J.F., see Witter, M.P., 164
Struble, R.G., see Brilliant, M., 23

Takahashi, K., Kameda, H., Kataoka, M., Ueno, S. and Akaike, N., Effects of Ca^{2+} antagonists and antiepileptics on tetrodotoxin-sensitive Ca^{2+}-conducting channels in isolated rat hippocampal CA1 neurons, 60
Takamatsu, M., see Ishiguro, K., 202
Takazawa, K., see Mailleux, P., 177
Temesvári, P., see Megyeri, P., 137
Tomura, H., Okajima, F. and Kondo, Y., Enkephalin induces Ca^{2+} mobilization in single cells of bradykinin-sensitized differentiated neuroblastoma hybridoma (NG108-15) cells, 93
Torres, G., Bitran, M. and Huidobro-Toro, J.P., Co-release of neuropeptide Y (NPY) and noradrenaline from the sympathetic nerve terminals supplying the rat vas deferens; influence of calcium and the stimulation intensity, 39
Toyota, T., see Shimosegawa, T., 67
Trifaro, J.-M., see Geertsen, S., 207

Uchida, T., see Ishiguro, K., 202
Ueno, S., see Takahashi, K., 60
Ungerstedt, U., see Morino, P., 133

Valtschanoff, J.G., Weinberg, R.J., Rustioni, A. and Schmidt, H.H.H.W., Nitric oxide synthase and GABA colocalize in lamina II of rat spinal cord, 6
Vanderhaeghen, J.-J., see Mailleux, P., 173, 177
Vas, T., see Megyeri, P., 137
Vázquez, R., see Alonso, J.R., 101
Völker, A., see Ballanyi, K., 141

Wakizaka, A., see Fujii, S., 148
Watt, C.B. and Florack, V.J., A double-label analysis demonstrating the non-coexistence of tyrosine hydroxylase-like and GABA-like immunoreactivities in amacrine cells of the larval tiger salamander retina, 47
Webb, C.L., see Lysko, P.G., 34
Weider, J.M., see Luebke, J.I., 129
Weinberg, R.J., see Valtschanoff, J.G., 6
Wilcox, G.L., see Kitto, K.F., 1
Willis, W.D., see Paleckova, V., 19
Witter, M.P., Daelmans, H.E.M., Jorritsma-Byham, B., Staiger, J.F. and Wouterlood, F.G., Restricted origin and distribution of projections from the lateral to the medial septal complex in rat and guinea pig, 164
Wouterlood, F.G., see Witter, M.P., 164

Yoshida, K., see Shimosegawa, T., 67